



MS-A0211 / Period III 2020

Final Exam, 17.02.2020

Aalto University

No calculators or notes of any kind are allowed.

This exam consists of 6 problems, each of equal weight.

Notation for vectors: $\langle a, b, c \rangle = ai + bj + ck$.

Spherical coordinates: $x = \rho \sin(\phi) \cos(\theta)$, $y = \rho \sin(\phi) \sin(\theta)$, $z = \rho \cos(\phi)$ and " $dV = \rho^2 \sin(\phi)$ ".

Question 1: Here are some unrelated direct questions

- (a) Consider the limits $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x + y}$, $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$, and $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{e^{x+y}}$.

Find a limit that does not exist and justify why it does not exist (you do NOT have to make any comment about limits that do exist).

- (b) Let $\mathbf{r}(t) = \langle t^2, t \rangle$ for $0 \leq t \leq 2$. Sketch the curve and write an integral (purely in terms of t) for the arc length of this curve. You do NOT have to evaluate the integral.
- (c) Compute the double integral of $f(x, y) = x^2y$ over the triangular region with vertices $(0, 0)$, $(-1, 1)$, and $(1, 1)$.

Question 2: Let $f(x, y) = \sin(xe^y) - x + 3$. A given fact is that $f(0.2, 0.1) = 3.01924$ accurate to 5 decimal places.

- (a) Compute all the 1st and 2nd order partial derivatives of $f(x, y)$
- (b) Taking $(0, 0)$ as the reference point, use *linear approximation* to find an approximation of $f(0.2, 0.1)$.
- (c) By referring to the idea of a tangent plane, explain why you obtained the answer you did in part (b)
- (d) Use a *2nd order Taylor polynomial* to find an approximation of $f(0.2, 0.1)$. Is this approximation better or worse than the linear approximation?
- (e) Find a critical point of $f(x, y)$ and determine if it is a local min, local max or saddle.
-

Question 3: Let $f(x, y) = 4xy$

- (a) Find the absolute extrema (i.e., min and max) of $f(x, y)$ on the elliptical region $4x^2 + y^2 \leq 8$. Determine all the locations of the extreme values.
- (b) Does $f(x, y)$ have an absolute minimum and absolute maximum on the 1st quadrant (that is, the region $\{(x, y) \mid x \geq 0, y \geq 0\}$). Justify your answer.

Question 4: Let $f(x, y) = y - x^2$.

- (a) Write the equations and then make a large sketch of the level curves $f(x, y) = 1$ and $f(x, y) = 2$ on the same axes (that is, make a contour plot with these two level curves).
- (b) On the above plot, draw a point P and a vector \mathbf{u} (with initial point at P) such that the directional derivative $D_{\mathbf{u}}f(P)$ is negative.
- (c) At the point $(1, 3)$, find the direction (vector) in which $f(x, y)$ is increasing the fastest. Sketch the point and the vector on the above plot. Does it look approximately correct? Why or why not?

Question 5: Two double integral questions.

- (a) Reverse the order of integration for $\int_0^1 \int_0^{2x^2+1} f(x, y) dy dx$. That is write as an integral of the form $\iint \dots dx dy$.
- (b) Compute the double integral of the function $f(x, y) = e^{x^2+y^2}$ over the top half of the disk of radius 3 centered at $(0, 0)$.

Question 6: Two triple integral questions.

- (a) Set up an integral in *cylindrical coordinates* to represent the volume of the region in the first octant (i.e. $x \geq 0, y \geq 0, z \geq 0$) that lies above the xy -plane, below the plane $z = y - x$ and inside the cylinder $x^2 + y^2 = 4$. Do NOT evaluate the integral.
- (b) Let E be the solid region that lies inside the sphere $x^2 + y^2 + z^2 = 2$ and above the plane $z = 1$. Let $d(x, y, z) = z^2$ be the density of E . Sketch E and write a triple integral in *spherical coordinates* that gives the mass of E . You do NOT have to evaluate the integral.